



Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : [http://oatao.univ-toulouse.fr/Eprints ID : 4353](http://oatao.univ-toulouse.fr/Eprints/ID/4353)

To cite this version :

TROEGELER, Annabelle, NICOT M C, ENJALBERT, Francis.
Enzymatic approach of linoleic acid ruminal biohydrogenation. *XIe Symposium International sur la Physiologie des Ruminants*. 6-9
Sept 2009. Clermont-Ferrand, France : 2009.


Any correspondance concerning this service should be sent to the repository administrator: staff-oatao@inp-toulouse.fr.

Enzymatic approach of linoleic acid ruminal biohydrogenation




A. Troegeler-Meynadier^{1,2*}, M.C. Nicot^{1,2} and F. Enjalbert^{1,2}
¹Université de Toulouse ; INPT, ENVT ; UMR 1289 Tandem, F-31076 Toulouse, France
²INRA ; UMR 1289 Tandem, F-31326 Castanet-Tolosan, France
a.troegeler@envt.fr

Ruminal biohydrogenation (BH) corresponds to a microbial reduction of dietary unsaturated fatty acid (Fig. 1). The control of BH reactions is of interest for researchers because BH directly affects the composition of fatty acids of milk and meat. In order to better understand C18:2 BH and its variations, the development of an enzymatic approach is necessary to ascertain if the action of modulators affects the bacterial enzyme activity or ruminal bacteria. The aim of this study was to investigate the C18:2 BH capacity of ruminal content after inactivation of bacteria by chloramphenicol, an inhibitor of protein synthesis in prokaryotes.



Materials and methods




Results and discussion

Table 1: Evolution of C18:0, C18:2, CLA and C18:1 isomers amounts (mg) during C18:2 BH.

Duration of incubation	0h	1h	2h	3h	SEM	P	
C18:0	0.567 ^a	0.581 ^a	0.584 ^a	0.615 ^b	0.07	0.006	→ C18:0 was only significantly (P<0.05) increased after 3h of incubation
<i>cis9</i> -C18:1	0.044	0.04	0.042	0.043	0.001	0.631	
<i>cis11</i> -C18:1	0.005	0.005	0.005	0.006	0.001	0.661	
<i>cis12</i> -C18:1	0.002 ^a	0.004	0.006	0.007 ^b	0.001	0.031	→ <i>cis12</i> -C18:1 and <i>trans6+7+8</i> -C18:1 weakly increased.
<i>cis15</i> -C18:1	0.001	0.001	0.000	0.000	0.000	0.131	
<i>trans6+7+8</i> -C18:1	0.004 ^a	0.004	0.005 ^b	0.005	0.000	0.013	
<i>trans9</i> -C18:1	0.002	0.002	0.002	0.002	0.000	0.936	
<i>trans10</i> -C18:1	0.003 ^a	0.000	0.025	0.032 ^b	0.006	0.047	→ <i>trans10</i> -C18:1 and <i>trans11</i> -C18:1 were the most abundant <i>trans</i> -C18:1 isomers, and increased throughout incubation
<i>trans11</i> -C18:1	0.056 ^a	0.087	0.105 ^b	0.115 ^b	0.007	0.002	
<i>trans12</i> -C18:1	0.009	0.007	0.005	0.007	0.001	0.560	
<i>trans13+14</i> -C18:1	0.000	0.000	0.000	0.000	0.000	-	
<i>trans15</i> -C18:1	0.004	0.004	0.004	0.004	0.000	0.712	
<i>trans16</i> -C18:1	0.007	0.006	0.006 ^a	0.007 ^b	0.000	0.036	
C18:2	1.074 ^a	0.749 ^b	0.592 ^{bc}	0.508 ^c	0.043	<0.001	→ C18:2 decreased during incubation
<i>trans10,cis12</i> -CLA	0.001 ^a	0.028 ^b	0.047 ^c	0.074 ^d	0.003	<0.001	→ <i>trans10,cis12</i> -CLA and <i>cis9,trans11</i> -CLA were the most abundant CLA isomers, and increased throughout incubation, except <i>cis9,trans11</i> -CLA max after 1h
<i>cis9,cis11</i> -CLA	0.000	0.000	0.000	0.000	0.000	-	
<i>cis9,trans11</i> -CLA	0.003 ^a	0.034 ^b	0.024	0.019	0.006	0.029	
<i>trans9,trans11</i> -CLA	0.000 ^a	0.004 ^b	0.005 ^c	0.008 ^d	0.000	<0.001	

Figure 2: evolution of intermediates of linoleic acid biohydrogenation



- BH of C18:2 \Rightarrow *cis9,trans11*-CLA + *trans10,cis12*-CLA \Leftrightarrow *trans11*-C18:1 and *trans10*-C18:1
- *cis12*-C18:1 \Leftrightarrow *trans10,cis12*-CLA,
- *trans6+7+8*-C18:1 from the reduction of minor CLA isomers not quantified in this study (Shingfield *et al.*, 2008).
- *trans11* pathway was rapid compared to *trans10* pathway which was slow:

 1. *trans10,cis12*-CLA accumulated vs. *cis9,trans11*-CLA max at 1h \Rightarrow after 3h: *trans10,cis12*-CLA > *cis9,trans11*-CLA
 2. *trans10*-C18:1 increased after 2h and < *trans10,cis12*-CLA.

- C18:0 began to increase in the media when *trans11*-C18:1 concentration was over 0.05 mg/mL.

Conclusion : Such evolution of fatty acids involved in C18:2 BH was similar to that reported in vitro with living ruminal microorganisms by Harfoot *et al.* (1973) and Jouany *et al.* (2007). This approach using Cm could be an interesting and valid method to study enzymes involved in C18:2 BH independently of bacteria, however 3h of incubation were not sufficient to study the final reduction.

Allison, J.L., R.E. Hartman, R.S. Hartman, A.D. Wolfe, J. Ciak and F.E. Hahn, 1962. Mode of action of chloramphenicol. J. Bacteriol. 83: 609-615.
 Harfoot, G.C., R.C. Noble and J.H. Moore, 1973. Factors influencing the extent of biohydrogenation of linoleic acid by rumen microorganisms in vitro. J. Sci. Food Agric. 24: 961-970.
 Jouany, J.P., B. Lassalas, M. Doreau and F. Glasser, 2007. Dynamic features of the rumen metabolism of linoleic acid, linolenic acid and linseed oil measured in vitro. Lipids. 42: 351-360.
 Rocha, E.R., T. Selby, J.P. Coleman and C.J. Smith, 1996. Oxidative stress response in an anaerobe, Bacteroides fragilis: a role for catalase in protection against hydrogen peroxide. J. Bacteriol. 178: 6895-6903.
 Shingfield, K.J., S. Ahvenjärvi, V. Toivonen, A. Vanhatalo, P. Huhtanen and J. M. Griinari, 2008. Effect of incremental levels of sunflower-seed oil in the diet on ruminal lipid metabolism in lactating cows. Br. J. Nutr. 99: 971-983.