



OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>
Eprints ID: 8053

To cite this document: Fontane, Jérôme and Dritschel, David and Scott, Richard
On the validity of Kraichnan scalings for forced two-dimensional turbulence.
(2012) In: 65th Annual Meeting of the American Physical Society's Division of
Fluid Dynamics (DFD), 18-20 Nov 2012, San Diego, United States.
(Unpublished)

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@inp-toulouse.fr

On the validity of Kraichnan scalings for forced two-dimensional turbulence

Jérôme Fontane¹, David G. Dritschel² and Richard K. Scott²

¹*Université de Toulouse, ISAE, 10 Av. Édouard Belin, 31055 Toulouse, France*

²*School of Mathematics and Statistics, University of St Andrews, St Andrews KY16 9SS, United Kingdom*

We examine the validity of the scaling laws derived by Kraichnan (1967) for forced two-dimensional turbulence. We use a new numerical technique (Dritschel & Fontane 2010) to reach higher Reynolds number than previously accessible with classical pseudo-spectral methods. No large scale friction or hypo-diffusion is used in order to avoid any distortion of the inverse cascade and to be in agreement with the theoretical framework used by Kraichnan. Both spectral and spatial forcing are considered and statistical convergence is obtained through large simulation ensembles.

A steeper energy spectrum proportional to k^{-2} is observed for the inverse cascade in place of the classical $k^{-5/3}$ prediction. This steepening is shown to be associated with a faster growth of energy at large scales, scaling like t^{-1} rather than Kraichnan's prediction of $t^{-3/2}$. The deviation from Kraichnan's theory is related to the emergence of a vortex population dominating the distribution of energy across scales, and whose number density and vorticity distribution with respect to vortex area are related to the shape of the enstrophy spectrum.

References

- DRITSCHEL, D. G. & FONTANE, J. 2010 The combined Lagrangian advection method. *J. Comp. Phys.* **229**, 5408–5417.
- KRAICHNAN, R. 1967 Inertial ranges in two-dimensional turbulence. *Phys. Fluids.* **10**, 1417–1423.